

Fitodenta hydrogel with povidone-iodine (PVP-I)

Clinical use protocol for dental professionals

This document provides dental professionals with practical guidance on using Fitodenta hydrogel with povidone-iodine (PVP-I) after dental procedures to ensure appropriate wound antisepsis, hygiene and postoperative care.

1. Product description

Fitodenta hydrogel with povidone-iodine (PVP-I) is a topical antiseptic hydrogel intended for postoperative wound care, disinfection and antiseptic action.

The product combines the antiseptic properties of povidone-iodine with hyaluronic acid, which helps maintain moisture balance and tissue comfort.

2. Indications

- After tooth extraction (antisepsis of the extraction socket)
- After oral surgical procedures (implant placement, soft-tissue surgery)
- For wounds or mucosal lesions in the oral cavity
- When additional local antiseptic care is needed

3. Recommendations for use

- After completing the procedure, gently apply a small amount of hydrogel directly to the wound or operated area.
- Reapplication 2–3 times a day is recommended during the first few days after the procedure.
- Adjust the duration of use according to the clinical situation and course of healing.

4. Properties and action

- Povidone-iodine has broad-spectrum antiseptic activity against bacteria, viruses and fungi.
- Hyaluronic acid helps maintain appropriate moisture balance and comfort during healing.

5. Safety information

- For local use in the oral cavity only.
- Avoid contact with the eyes.
- Do not use in patients allergic to iodine or other ingredients.
- Discontinue use if adverse reactions occur.

6. Integration into the treatment plan

- May be used as an additional antiseptic measure after dental procedures.
- Helps maintain a clean wound environment during healing.

Note: This preparation is intended for professional use in dental practice.